

Table B. Percent of fetal death records on which specified items were not stated: Each State, 1999  
 [By place of occurrence. Records include only those with stated or presumed period of gestation of 20 weeks or more.]

Area	Length of gestation 2/	Marital status	Place of delivery	Birth weight	Month prenatal care began	Number of prenatal visits	Hispanic origin of mother
Total of reporting areas 1	3.4	34.4	0.0	11.2	16.0	17.3	6.0
Alabama	0.2	0.7	-	3.8	4.1	5.1	1.5
Alaska	-	-	-	8.2	-	2.0	2.0
Arizona	1.8	0.4	-	6.3	9.5	12.5	2.6
Arkansas	0.4	-	-	4.7	9.8	8.1	-
California	10.8	- - -	0.0	1.3	5.1	6.7	1.4
Colorado	0.7	5.3	-	12.0	20.6	19.9	11.3
Connecticut	-	0.4	-	4.7	24.2	25.8	7.2
Delaware	3.9	-	-	3.9	7.8	15.7	2.0
District of Columbia	1.7	12.0	-	1.7	25.6	29.1	2.6
Florida	2.4	1.9	-	8.8	19.7	19.6	4.5
Georgia	0.8	0.4	-	17.0	24.5	25.6	19.0
Hawaii	0.8	5.5	0.8	46.9	40.6	36.7	14.1
Idaho	0.9	-	-	7.3	7.3	6.4	-
Illinois	1.4	5.7	-	8.7	14.9	15.8	3.5
Indiana	1.2	2.3	-	13.8	11.3	12.3	6.6
Iowa	-	0.5	-	2.7	4.1	7.2	-
Kansas	1.7	-	-	-	3.4	9.0	3.9
Kentucky	1.2	0.3	-	4.8	4.2	5.1	-
Louisiana	2.0	2.2	-	5.5	6.7	8.3	0.8
Maine	7.5	19.4	-	38.8	19.4	28.4	16.4
Maryland	2.4	2.1	0.3	5.2	9.1	13.1	4.6
Massachusetts	0.7	1.6	-	5.8	16.7	14.6	22.0
Michigan	1.1	- - -	0.1	1.8	13.8	17.2	14.1
Minnesota	15.5	19.9	-	33.6	27.4	25.6	4.1
Mississippi	0.9	-	-	1.9	5.7	19.8	0.2
Missouri	0.4	0.2	-	1.5	7.2	9.2	1.1
Montana	-	1.6	-	4.8	-	7.9	-
Nebraska	-	1.4	-	2.7	2.7	2.0	4.1
Nevada	1.8	- - -	-	21.6	20.5	22.8	1.8
New Hampshire	-	-	-	5.9	1.5	4.4	17.6
New Jersey	3.2	4.2	-	20.5	17.2	19.4	2.0
New Mexico	-	-	-	1.1	1.1	5.3	3.2
New York State	1.8	- - -	0.1	37.9	28.9	27.8	2.8
New York City	1.2	- - -	-	28.2	44.1	42.4	9.5
North Carolina	0.9	0.2	-	7.0	8.4	8.5	0.4
North Dakota	-	-	-	4.5	4.5	4.5	11.4

Table B. Percent of fetal death records on which specified items were not stated: Each State, 1999 - cont.  
 [By place of occurrence. Records include only those with stated or presumed period of gestation of 20 weeks or more.]

	Length of gestation 2/	Marital status	Place of delivery	Birth weight	Month prenatal care began	Number of prenatal visits	Hispanic origin of mother
Ohio	0.7	25.2	-	6.6	4.8	6.8	0.9
Oklahoma	57.3	41.4	0.7	39.1	56.0	54.6	- - -
Oregon	0.4	-	-	0.8	2.4	2.4	1.2
Pennsylvania	2.3	4.3	-	8.4	14.1	15.3	2.5
Rhode Island	-	90.8	-	22.4	98.7	97.4	96.1
South Carolina	0.4	0.2	-	2.9	5.5	5.9	1.8
South Dakota	-	-	-	-	4.8	4.8	-
Tennessee	0.9	-	-	0.7	7.2	7.4	4.7
Texas	4.5	- - -	-	9.7	15.3	17.5	1.6
Utah	-	-	-	9.6	12.8	13.7	-
Vermont	-	-	-	-	10.3	3.4	3.4
Virginia	0.5	3.9	-	37.8	40.9	43.3	9.3
Washington	2.6	3.2	-	15.3	21.6	25.0	7.3
West Virginia	1.3	-	-	4.6	9.8	11.8	-
Wisconsin	0.6	-	-	1.4	1.1	1.4	-
Wyoming	-	-	-	-	7.7	3.8	-

Table B. Percent of fetal death records on which specified items were not stated: Each State, 1999 - cont.  
 [By place of occurrence. Records include only those with stated or presumed period of gestation of 20 weeks or more.]

	Mother's educational attainment	Medical risk factors 3/	Tobacco use	Alcohol use	Obstetric procedures	Complication of labor and/ or delivery 4/	Congenital anomalies
Total of reporting areas 1	15.7	10.3	24.0	22.3	7.6	9.2	16.0
Alabama	5.5	0.9	2.6	2.4	1.4	2.6	3.6
Alaska	8.2	-	12.2	16.3	8.2	12.2	28.6
Arizona	2.8	1.6	5.3	6.3	3.6	1.8	3.0
Arkansas	12.3	1.7	1.3	1.3	0.9	1.7	0.9
California	6.2	4.0	---	---	2.7	3.4	4.3
Colorado	21.1	13.2	15.3	16.2	7.4	13.0	11.3
Connecticut	43.6	11.4	16.1	19.5	7.6	15.7	27.5
Delaware	5.9	9.8	17.6	17.6	9.8	15.7	15.7
District of Columbia	31.6	51.3	59.8	63.2	53.0	54.7	63.2
Florida	14.0	3.1	8.1	8.8	2.6	3.6	4.0
Georgia	36.3	4.6	12.2	12.5	3.7	4.8	4.7
Hawaii	30.5	---	---	---	---	---	---
Idaho	6.4	0.9	3.6	3.6	1.8	2.7	3.6
Illinois	10.1	18.2	16.8	19.9	13.4	17.9	21.6
Indiana	9.9	9.0	9.4	10.5	6.8	7.0	10.9
Iowa	1.8	1.4	0.9	1.4	0.5	1.8	3.2
Kansas	5.1	16.9	16.9	16.9	10.7	8.4	17.4
Kentucky	2.4	26.6	23.3	23.9	24.2	26.0	35.5
Louisiana	9.1	2.0	6.9	5.9	1.8	2.2	2.2
Maine	14.9	22.4	14.9	20.9	16.4	16.4	28.4
Maryland	14.6	6.2	-	-	4.6	5.1	5.9
Massachusetts	20.0	14.8	9.3	10.4	8.1	8.8	17.2
Michigan	24.5	0.9	5.6	6.9	0.6	1.0	0.6
Minnesota	10.3	18.1	14.2	15.5	9.6	15.8	17.3
Mississippi	7.1	2.1	4.2	4.2	0.7	1.9	3.1
Missouri	9.8	1.1	3.9	4.4	2.0	1.5	2.6
Montana	1.6	7.9	11.1	11.1	4.8	7.9	6.3
Nebraska	2.7	3.4	4.8	6.1	2.0	2.0	4.8
Nevada	5.3	27.5	36.3	38.6	24.6	34.5	29.8
New Hampshire	11.8	-	2.9	1.5	-	-	-
New Jersey	16.4	10.0	6.7	7.5	6.2	11.6	12.0
New Mexico	20.2	1.1	1.1	2.1	1.1	-	---
New York State	42.2	27.2	---	18.9	10.1	17.3	40.1
New York City	32.3	22.0	23.8	25.2	21.7	24.1	---
North Carolina	4.6	0.1	3.4	4.7	-	-	-
North Dakota	9.1	6.8	2.3	9.1	6.8	9.1	11.4

Table B. Percent of fetal death records on which specified items were not stated: Each State, 1999 - cont.  
 [By place of occurrence. Records include only those with stated or presumed period of gestation of 20 weeks or more.]

	Mother's educational attainment	Medical risk factors 3/	Tobacco use	Alcohol use	Obstetric procedures	Complication of labor and/ or delivery 4/	Congenital anomalies
Ohio	3.9	3.6	6.2	8.2	3.1	4.5	7.5
Oklahoma	45.0	---	---	---	---	---	---
Oregon	7.5	-	2.4	2.4	-	-	-
Pennsylvania	18.5	2.8	6.8	8.6	2.3	3.2	6.9
Rhode Island	96.1	73.7	80.3	82.9	73.7	76.3	82.9
South Carolina	7.8	1.8	3.3	4.3	2.2	1.6	3.9
South Dakota	-	-	---	---	-	-	-
Tennessee	13.4	4.0	5.6	6.0	3.6	5.1	8.3
Texas	7.5	14.5	2.5	3.1	1.5	2.5	0.1
Utah	6.4	1.4	2.7	2.3	-	3.7	-
Vermont	6.9	6.9	-	-	6.9	-	10.3
Virginia	42.2	9.4	9.4	9.4	9.4	9.4	9.4
Washington	17.2	17.0	15.5	32.5	15.7	17.5	25.0
West Virginia	11.8	0.7	11.1	15.0	0.7	0.7	-
Wisconsin	2.9	0.6	-	-	-	0.3	4.9
Wyoming	3.8	-	7.7	11.5	-	-	-

0.0 Quantity more than zero but less than 0.05.

--- Data not available

- Quantity zero

1/ Excludes data for Puerto Rico, Virgin Islands, Guam, American Samoa, and the Commonwealth of the Northern Marianas.

2/ California and Oklahoma report date last normal menses began but do not report clinical estimate of gestation.

3/ Kansas and South Dakota do not report Rh sensitization; Texas does not report genital herpes and uterine bleeding.

4/ Texas does not report cephalopelvic disproportion, anesthetic complications, and fetal distress.